

Euler Summation Formula for Path Integrals

A. Bogojević, A. Balaž, and A. Belić
Institute of Physics, Belgrade, Serbia and Montenegro

Abstract

We present and comment on some details of a new analytical method [1, 2] for systematic improvement of the convergence of path integrals of a generic N -fold discretized theory. The new method represents an Euler summation formula for path integrals. Keeping the first p terms in this formula improves convergence of path integrals to the continuum limit to $1/N^p$. We have given explicit calculations up to $p = 9$. We have also checked this derived speedup in convergence by performing Monte Carlo simulations on several different models.

1 Introduction

Feynman's path integrals have, since their inception [3, 4], represented an extremely compact and rich formalism for dealing with quantum theories. They have been powerful tools for dealing with symmetries (including gauge symmetry), for deriving non-perturbative results (such as solitons, instantons, symmetry breaking), for showing connections between different theories or different sectors of theories (e.g. bosonization, duality) [5, 6]. Their flexibility and intuitive appeal have allowed us to generalize quantization to ever more complicated systems. As a result, path integrals have led to a rich cross fertilization of ideas between high energy and condensed matter physics [7, 8]. Today, they are used both analytically and numerically [9, 10, 11, 12, 13] in many other areas of physics, chemistry and materials science. They are starting to play more prominent roles in several areas of mathematics and in modern finance [14]. An extensive review of path integrals and their applications can be found in [15].

This was the good news. The bad news is that we still have very little knowledge of the precise mathematical properties of path integrals. In addition, a very small number of path integrals can be solved exactly. Although the functional formalism has been used to derive many general approximation techniques (perturbation, semi-classical expansion, variational methods) along with a host of model-specific approximations, there still remain many interesting models that can't be analyzed analytically and need to be treated numerically. The definition of path integrals as a limit of multiple integrals makes their numerical evaluation quite natural. The most all-around applicable numerical method for such calculations is based on Monte Carlo simulations. However, numerical integration of path integrals is notoriously demanding of computing time. In fact specific path integral calculations serve as benchmarks for new generations of supercomputers.

Several research groups have in the past focused on improving the convergence of path integrals. The best available result for a generic theory (valid

only for partition functions and not for general amplitudes) is the convergence of N -fold discretized expressions as $1/N^4$ [16, 17, 18]. Related investigations have focused either on improvements in short time propagation [19, 20, 21] or have presented model specific improvements of the action [22, 23]. Ref. [24] gives a useful comparison of several different approaches.

In order to further significantly speed up numerical procedures for calculating path integrals for a generic theory it is necessary to add new analytical input. We present and comment on the systematic investigation of the relation between different discretizations of a given theory derived in [1, 2]. A result of this investigation is a procedure for constructing a series of effective actions $S^{(p)}$ having the same continuum limit as the starting action S , but which approach that limit as $1/N^p$. Using this procedure we have obtained explicit expressions for these effective actions for $p \leq 9$. In the current paper we cast the new analytical input in the form of a Euler summation formula for path integrals. The fact that it is possible to derive such a formula for path integrals of a generic theory hints at the possible existence of (currently unseen) simplifications that might make it possible to set up a rigorous theory of path integration – a modern day parallel to Riemann’s integration theory.

2 Ordinary Integrals

The current status of the development of the path integral formalism is quite similar to that of ordinary integrals before the setting up of integration theory by Riemann. In those days integrals were calculated directly from the defining formula, i.e. one looked at a specific discretization of the integral (Darboux sum), attempted to do the sum explicitly, and finally tried to calculate the continuum limit. For example,

$$I[f] \equiv \int_0^T f(t)dt = \lim_{N \rightarrow \infty} I_N[f] , \quad (1)$$

where

$$I_N[f] = \sum_{n=1}^N f(t_n)\epsilon_N , \quad (2)$$

$\epsilon_N = T/N$ and $t_n = n\epsilon_N$. It goes without saying that done this way, even the simplest ordinary integrals presented a challenge. The mathematicians of the 18th century did not have computers at their disposal or the development of integration theory might have come much later, i.e. they might have succumbed to doing brute force numerical calculations of integrals of all but the simplest functions. The problem with these hypothetical numerical calculations would have been two fold: they would have been inefficient (the discretized sums converge slowly to the continuum value), and they would have worked (thus quite probably slowing down the further development of integration theory). Luckily, this early numerical road was not open. The last great step in the development of integration before Riemann was made by Euler. Let us look at a simple derivation of Euler’s summation formula that we will generalize to path integrals in the following sections.

Discretization is not unique. This makes it possible to change $f(t)$ to some other function (adding terms proportional to ϵ_N , ϵ_N^2 , etc.) without changing

the integral. Let us assume that $f^*(t)$ is such an equivalent function with the added property that the sums $I_N[f^*]$ do not depend on N . In fact we shall present a way of explicitly constructing $f^*(t)$ for any given $f(t)$. We first look at the simple case of $f(t) = 1$. Now

$$I_N[1] = \sum_{n=1}^N \epsilon_N = T, \quad (3)$$

which is already N -independent. Hence, in this case, all the additional terms vanish. Note that $f^*(t)$ is completely determined by the original function $f(t)$ (and by ϵ_N), so that the additional terms necessarily depend only on the derivatives f' , f'' , etc.

The second step is to take $f(t) = t$. In this case we get

$$I_N[t] = \sum_{n=1}^N t_n \epsilon_N = \frac{N(N+1)}{2} \frac{T^2}{N^2} = \frac{T^2}{2} + \frac{T^2}{2N}. \quad (4)$$

From this it follows that $I_N[t - \frac{\epsilon_N}{2}] = \frac{T^2}{2}$. Therefore, up to f'' and higher derivatives of f that all vanish for linear $f(t)$, we have $f^*(t) = f(t) - \frac{\epsilon_N}{2} f'(t)$.

We continue this procedure by looking at $f(t) = t^2$. In this case we find

$$I_N[t^2] = \sum_{n=1}^N t_n^2 \epsilon_N = \frac{N(N+1)(2N+1)}{6} \frac{T^3}{N^3} = \frac{T^3}{3} + \frac{T^3}{2N} + \frac{T^3}{6N^2}. \quad (5)$$

It follows that $I_N[t^2 - \epsilon_N t_n - \frac{2}{3} \epsilon_N^2] = \frac{T^3}{3}$. In terms of f^* this gives $f^*(t) = f(t) - \frac{\epsilon_N}{2} f'(t) - \frac{2\epsilon_N^2}{3} f''(t) + \dots$. The additional terms now depend on higher powers of ϵ_N as well as on higher derivatives and are determined by considering $I_N[t^3]$, and so on. In this way we have constructed a procedure for finding $f^*(t)$ for any given $f(t)$. Remembering that $I_N[f^*]$ does not depend on N we find

$$\int_0^T f(t) dt = \sum_{n=1}^N f(t_n) \epsilon_N - \frac{\epsilon_N}{2} \sum_{n=1}^N f'(t_n) \epsilon_N - \frac{2\epsilon_N^2}{3} \sum_{n=1}^N f''(t_n) \epsilon_N + \dots \quad (6)$$

This is the well-known Euler summation formula. We may also write it more compactly as

$$I[f] = I_N[f^{(p)}] + O(\epsilon_N^p), \quad (7)$$

where $f^{(p)}$ is the truncation of f^* to the first p terms. The Euler formula gives the analytical relation between integrals and their discretized sums. Looked at numerically, this formula allows us to increase the speed of convergence of discretized expressions to the continuum limit. For example, in the defining relation the discretized expressions differ from the continuum by a term of order $O(1/N)$. By using the Euler sum formula with p terms we can reduce that error to $O(1/N^p)$. All that is needed to do this is that the integrand is differentiable $p-1$ times. In the following sections we will generalize the above approach to path integrals.

3 General Properties of Path Integrals

In the functional formalism the quantum mechanical amplitude $A(a, b; T) = \langle b | e^{-T\hat{H}} | a \rangle$ is given in terms of a path integral which is simply the $N \rightarrow \infty$ limit of the $(N - 1)$ - fold integral expression

$$A_N(a, b; T) = \left(\frac{1}{2\pi\epsilon_N} \right)^{\frac{N}{2}} \int dq_1 \cdots dq_{N-1} e^{-S_N} . \quad (8)$$

The Euclidean time interval $[0, T]$ has been subdivided into N equal time steps of length $\epsilon_N = T/N$, with $q_0 = a$ and $q_N = b$. S_N is the naively discretized action of the theory. We focus on actions of the form

$$S = \int_0^T dt \left(\frac{1}{2} \dot{q}^2 + V(q) \right) , \quad (9)$$

whose naive discretization is simply

$$S_N = \sum_{n=0}^{N-1} \left(\frac{\delta_n^2}{2\epsilon_N} + \epsilon_N V_n \right) , \quad (10)$$

where $\delta_n = q_{n+1} - q_n$, $V_n = V(\bar{q}_n)$, and $\bar{q}_n = \frac{1}{2}(q_{n+1} + q_n)$. We use units in which \hbar and particle mass equal 1.

As was the case with ordinary integrals the definition of the path integrals also makes it necessary to make the transition from the continuum to the discretized theory, a process that is far from unique. For theories described by eq. (9) we have the freedom to choose any point in $[q_n, q_{n+1}]$ in which to evaluate the potential without changing physics – the discretized amplitudes do differ, but they tend to the same continuum limit. The calculations we present turn out to be simplest in the mid-point prescription where the potential V is evaluated at \bar{q}_n . A more important freedom related to our choice of discretized action has to do with the possibility of introducing additional terms that explicitly vanish in the continuum limit. Actions with such additional terms will be called effective. For example, the term $\sum_{n=0}^{N-1} \epsilon_N \delta_n^2 g(\bar{q}_n)$, where g is regular when $\epsilon_N \rightarrow 0$, does not change the continuum physics since it goes over into $\epsilon_N^2 \int_0^T dt \dot{q}^2 g(q)$, i.e. it vanishes as ϵ_N^2 . Such terms do not change the physics, but they do affect the speed of convergence. A systematic study of the relation between different discretizations of the same path integral will allow us to explicitly construct a series of effective actions with progressively faster convergence to the continuum. Before we do this we will parallel the derivation in the previous section and derive some general properties of the best effective action.

The amplitude $A(a, b; T)$ of some theory with action S satisfies

$$A(a, b; T) = \int dq_1 \cdots dq_{n-1} A(b, q_{n-1}; \epsilon_N) \cdots A(q_1, a; \epsilon_N) , \quad (11)$$

for all N . This general relation is a direct consequence of the linearity of states in a quantum theory. In analogy with ordinary integrals let us now suppose that there exists an effective action S^* that is equivalent to S (i.e. that leads to the same continuum limit for all path integrals) with the additional property that

its N -fold discretized amplitude $A_N^*(a, b; T)$ does not depend on N , i.e. that satisfies

$$A_N^*(a, b; T) = A(a, b; T) . \quad (12)$$

As was the case in the previous section we will in fact construct a general procedure for evaluating this effective action. For actions of the form given in eq. (9) we may write the amplitude as

$$A(q_{n+1}, q_n; \epsilon_N) = \left(\frac{1}{2\pi\epsilon_N} \right)^{\frac{1}{2}} \exp \left(-\frac{\delta_n^2}{2\epsilon_N} \right) \mathcal{A}(q_{n+1}, q_n; \epsilon_N) , \quad (13)$$

where the reduced amplitude $\mathcal{A} \rightarrow 1$ as $\epsilon_N \rightarrow 0$. Writing S_N^* as

$$S_N^* = \sum_{n=0}^{N-1} \left(\frac{\delta_n^2}{2\epsilon_N} + \epsilon_N W_n^* \right) , \quad (14)$$

and using eq. (8), (11) and (12) we find

$$\exp(-\epsilon_N W_n^*) = \mathcal{A}(q_{n+1}, q_n; \epsilon_N) . \quad (15)$$

Note that W_n^* is reminiscent of some effective potential, so it should depend on \bar{q}_n , however, from the above relation we see that it must also depend on δ_n . In addition, W^* also has an explicit dependence on the discrete time step ϵ_N , hence

$$W_n^* = W^*(\delta_n, \bar{q}_n; \epsilon_N) . \quad (16)$$

As we have seen, the above functional form is a direct consequence of the linearity of quantum theory. The equivalence of S and S^* implies that $W^* \rightarrow V(\bar{q})$ when ϵ_N and δ go to zero. The final general property of W^* follows from the reality of amplitudes in the Euclidean formalism. Using the hermiticity of the Hamiltonian we find $A(a, b; T) = A(a, b; T)^\dagger = \langle b | e^{-T\hat{H}} | a \rangle^\dagger = \langle a | e^{-T\hat{H}} | b \rangle = A(b, a; T)$. In terms of W^* this gives us

$$W^*(\delta_n, \bar{q}_n; \epsilon_N) = W^*(-\delta_n, \bar{q}_n; \epsilon_N) , \quad (17)$$

or, said another way, only even powers of δ_n are present in the expansion of W^* :

$$W^*(\delta_n, \bar{q}_n; \epsilon_N) = g_0(\bar{q}_n; \epsilon_N) + \delta_n^2 g_1(\bar{q}_n; \epsilon_N) + \delta_n^4 g_2(\bar{q}_n; \epsilon_N) + \dots . \quad (18)$$

All the functions g_k are regular in the $\epsilon \rightarrow 0$ limit. The link to the starting theory is now simply $g_0(\bar{q}_n; \epsilon_N) \rightarrow V(\bar{q}_n)$ as ϵ_N goes to zero. This concludes the general properties of W^* . The remaining properties will be analyzed in the following section by studying the relation of discretizations of different coarseness.

4 Relation Between Different Discretizations

We start by studying the relation between the $2N$ -fold and N -fold discretizations of the same theory. From eq. (8) we see that we can write the $2N$ -fold amplitude as an N -fold amplitude given in terms of a new action \tilde{S}_N determined by

$$e^{-\tilde{S}_N} = \left(\frac{2}{\pi\epsilon_N} \right)^{\frac{N}{2}} \int dx_1 \dots dx_N e^{-S_{2N}} , \quad (19)$$

where S_{2N} is the $2N$ -fold discretization of the starting action. We have written the $2N$ -fold discretized coordinates Q_0, Q_1, \dots, Q_{2N} in terms of q 's and x 's in the following way: $Q_{2k} = q_k$ and $Q_{2k-1} = x_k$. Note that we have $q_0 = a$, $q_N = b$, while the $N - 1$ remaining q 's play the role of the dynamical coordinates in the N -fold discretized theory. The x 's are the N remaining intermediate points that we integrate over in eq. (19). It is not difficult to see that if we use the naively discretized action S_N one obtains for \tilde{S}_N an expression that is not of the same form as S_N .

Having in mind the results of the previous section it is best to use the effective action

$$S_N^* = \sum_{n=0}^{N-1} \left(\frac{\delta_n^2}{2\epsilon_N} + \epsilon_N W^*(\delta_n, \bar{q}_n; \epsilon_N) \right), \quad (20)$$

which gives the same result for both the $2N$ -fold and N -fold discretizations. Therefore, in this case we get

$$e^{-S_N^*} = \left(\frac{2}{\pi\epsilon_N} \right)^{\frac{N}{2}} \int dx_1 \dots dx_N e^{-S_{2N}^*}. \quad (21)$$

From this one easily finds

$$\begin{aligned} \exp[-\epsilon_N W^*(\delta_n, \bar{q}_n; \epsilon_N)] &= \\ &= \left(\frac{2}{\pi\epsilon_N} \right)^{\frac{1}{2}} \int_{-\infty}^{+\infty} dy \exp\left(-\frac{2}{\epsilon_N} y^2\right) F\left(\bar{q}_n + y; \frac{\epsilon_N}{2}\right), \end{aligned} \quad (22)$$

where

$$\begin{aligned} -\frac{2}{\epsilon_N} \ln F(x; \epsilon_N) &= g_0\left(\frac{q_{n+1} + x}{2}; \epsilon_N\right) + g_0\left(\frac{x + q_n}{2}; \epsilon_N\right) + \\ &+ (q_{n+1} - x)^2 g_1\left(\frac{q_{n+1} + x}{2}; \epsilon_N\right) + (x - q_n)^2 g_1\left(\frac{x + q_n}{2}; \epsilon_N\right) + \dots \end{aligned} \quad (23)$$

The above integral equation can be solved for the simple cases of a free particle and a harmonic oscillator, and gives the well known results. Note however that for a general case the integral in eq. (22) is in a form that is ideal for an asymptotic expansion [25]. The time step ϵ_N is playing the role of small parameter (in complete parallel to the role \hbar plays in standard semi-classical, or loop, expansion). As is usual, the above asymptotic expansion is carried through by first Taylor expanding $F\left(\bar{q}_n + y; \frac{\epsilon_N}{2}\right)$ around \bar{q}_n and then by doing the remaining Gaussian integrals. Assuming that $\epsilon_N < 1$ (i.e. $N > T$) we have

$$\begin{aligned} g_0(\bar{q}_n; \epsilon_N) + \delta_n^2 g_1(\bar{q}_n; \epsilon_N) + \delta_n^4 g_2(\bar{q}_n; \epsilon_N) + \dots &= \\ &= -\frac{1}{\epsilon_N} \ln \left[\sum_{m=0}^{\infty} \frac{F^{(2m)}\left(\bar{q}_n; \frac{\epsilon_N}{2}\right)}{(2m)!!} \left(\frac{\epsilon_N}{4}\right)^m \right]. \end{aligned} \quad (24)$$

Note that $F^{(2m)}(x; \epsilon_N)$ denotes the corresponding derivative with respect to x . All that remains is to calculate these expressions using eq. (23) and to expand all the g_k 's around the mid-point \bar{q}_n . This is a straight forward though tedious calculation. In this paper we will illustrate the general procedure for calculating

S^* by explicitly giving its expansion to order ϵ_N^3 :

$$\begin{aligned}
g_0(\bar{q}_n; \epsilon_N) &= g_0\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \epsilon_N \left[\frac{1}{4}g_1\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \frac{1}{32}g_0''\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) \right] + \\
&\quad + \epsilon_N^2 \left[\frac{3}{16}g_2\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) - \frac{1}{32}g_0'^2\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \frac{1}{2048}g_0^{(4)}\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \right. \\
&\quad \left. + \frac{3}{128}g_1''\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) \right] \\
g_1(\bar{q}_n; \epsilon_N) &= \frac{1}{4}g_1\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \frac{1}{32}g_0''\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \\
&\quad + \epsilon_N \left[\frac{3}{8}g_2\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \frac{1}{1024}g_0^{(4)}\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) - \frac{1}{64}g_1''\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) \right] \\
g_2(\bar{q}_n; \epsilon_N) &= \frac{1}{16}g_2\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \frac{1}{6144}g_0^{(4)}\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) + \frac{1}{128}g_1''\left(\bar{q}_n; \frac{\epsilon_N}{2}\right) .
\end{aligned} \tag{25}$$

In the above relations we expanded g_0 up to ϵ_N^2 , g_1 up to ϵ_N , etc. We also disregarded all the higher g_k 's. The reason for this is that the short time propagation of any theory satisfies $\delta_n^2 \propto \epsilon_N$ while the g_k term enters the action multiplied by δ_n^{2k} . In general, if we wish to expand the effective action to ϵ_N^p we need to evaluate only g_0 (up to ϵ_N^{p-1}) and the remaining $p-1$ functions g_k (up to ϵ_N^{p-1-k}). The task of calculating the effective action to large powers of ϵ_N is time-consuming and is best done with the help of a standard package for algebraic calculations such as Mathematica. Using Mathematica we determined the corresponding expressions for $p \leq 9$.

Although the above system of recursive relations is non-linear, it is in fact quite easy to solve if we remember that the system itself was derived via an expansion in ϵ_N . Having this in mind we first write all the functions as expansions in powers of ϵ_N that are appropriate to the level p we are working at. For $p=3$, we have

$$\begin{aligned}
g_0(\bar{q}_n; \epsilon_N) &= V(\bar{q}_n) + \epsilon_N R_1(\bar{q}_n) + \epsilon_N^2 R_2(\bar{q}_n) \\
g_1(\bar{q}_n; \epsilon_N) &= R_3(\bar{q}_n) + \epsilon_N R_4(\bar{q}_n) \\
g_2(\bar{q}_n; \epsilon_N) &= R_5(\bar{q}_n) .
\end{aligned} \tag{26}$$

Putting this into the Eq. (25) we determine the functions R_1 to R_5 in terms of V . The $p=3$ level solution equals

$$\begin{aligned}
g_0 &= V + \epsilon_N \frac{V''}{12} + \epsilon_N^2 \left[-\frac{V'^2}{24} + \frac{V^{(4)}}{240} \right] \\
g_1 &= \frac{V''}{24} + \epsilon_N \frac{V^{(4)}}{480} \\
g_2 &= \frac{V^{(4)}}{1920} .
\end{aligned} \tag{27}$$

Note that W^* depends only on the initial potential V and its derivatives (as well as on ϵ_N). One can similarly calculate the effective action S^* to any desired level p . We denote the p level truncation of the effective action as $S^{(p)}$. $S^{(p)}$ has the property that its N -fold amplitudes deviate from the continuum expressions as $O(\epsilon_N^p)$

$$A(a, b; T) = A_N^{(p)}(a, b; T) + O(\epsilon_N^p) . \tag{28}$$

Comparing this to eq. (7) we see that we have just derived the generalization of the Euler summation formula to path integrals. Just as with the ordinary Euler formula it gives the relation between path integrals and their discretizations to any given precision.

It is important to note that one solves for the effective action at level p but once for all theories, i.e. the solution that is found holds for all initial potentials. The only requirement for the level p solution is that the starting potential is differentiable $2p - 2$ times. Solutions for larger values of p are a bit more cumbersome, however, they are just as easy to use in simulations. We have found that the growth in complexity of the effective actions with increasing p has little effect on computation time for $p \leq 4$, while simulations with $p = 9$ are roughly ten times slower due to this. However, this is an extremely small price to pay for a gain of eight orders of magnitude in the speed of convergence. Expressions for effective actions up to $p = 9$ can be found on our web site [26].




Figure 1: Deviations from the continuum limit $|A_N^{(p)} - A|$ as functions of N for $p = 1, 2, 4$ and 6 for an anharmonic oscillator with quartic coupling $\lambda = 10$, time of propagation $T = 1$ from $a = 0$ to $b = 1$. N_{MC} was $9.2 \cdot 10^9$ for $p = 1, 2$, $9.2 \cdot 10^{10}$ for $p = 4$, and $3.68 \cdot 10^{11}$ for $p = 6$. Dashed lines correspond to appropriate $1/N$ polynomial fits to the data. Solid lines give the leading $1/N$ behavior. The level p curve has a $1/N^p$ leading behavior.

The analytical derivations presented work equally well in both the Euclidean and Minkowski formalism (with appropriate $i\epsilon$ regularization), i.e. they are directly applicable to quantum systems as well as to statistical ones. However, the Monte Carlo simulations used to numerically document our analytical results necessarily needed to be done in the Euclidean formalism. We analyzed in detail several models: the anharmonic oscillator with quartic coupling $V = \frac{1}{2} q^2 + \frac{\lambda}{4!} q^4$ and a particle moving in a modified Pöschl-Teller potential over a wide range

of parameters. In all cases we found agreement with eq. (28). Fig. 1 illustrates this behavior in the case of an anharmonic oscillator. We see that the p level data indeed differs from the continuum amplitudes as a polynomial starting with $1/N^p$. The deviations from the continuum limit $|A_N^{(p)} - A|$ become exceedingly small for larger values of p making it necessary to use ever larger values of N_{MC} so that the MC statistical error does not mask these extremely small deviations. For $p = 6$ we see that although we used an extremely large number of MC samples ($N_{MC} = 3.68 \cdot 10^{11}$) the statistical errors become of the same order as the deviations already at $N \gtrsim 8$. For $p = 9$ this is the case even for $N = 2$, i.e. we already get the continuum limit within a MC error of around 10^{-8} .

To conclude, we have presented an algorithm that leads to significant speedup of numerical procedures for calculating path integrals. The increase in speed results from new analytical input that comes from a systematic investigation of the relation between discretizations of different coarseness and that leads to a generalization of the Euler summation formula to path integrals. We have presented an explicit procedure for obtaining a set of effective actions $S^{(p)}$ that have the same continuum limit as the starting action S , but which approach that limit ever faster. Amplitudes calculated using the N -point discretized effective action $S_N^{(p)}$ satisfy $A_N^{(p)}(a, b; T) = A(a, b; T) + O(1/N^p)$. We have obtained and analyzed the effective actions for $p \leq 9$ and have documented the speedup up to $1/N^9$ by conducting Monte Carlo simulations of several different models. Several interesting properties of this procedure follow from the fact that the solutions were obtained using an asymptotic expansion. These additional properties will be presented and discussed in a future publication. Extension to $d > 1$ is also in progress. The derivation of higher dimensional analogues of integral equation (22) does not seem to present a problem. The asymptotic expansion used to solve it is also directly generalizable. However, the algebraic recursive relations that determine $S^{(p)}$ will be more complex and may practically limit us to smaller values of p .

Acknowledgement

We acknowledge financial support from the Ministry of Science and Environmental Protection of the Republic of Serbia through projects No. 1486 and No. 1899.

References

- [1] A. Bogojević, A. Balaž, and A. Belić, Phys. Rev. Lett. **94**, 180403 (2005).
- [2] A. Bogojević, A. Balaž, and A. Belić, Submitted to Phys. Rev. B
- [3] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
- [4] R. P. Feynman, *Statistical Mechanics* (W. A. Benjamin, New York, 1972).
- [5] C. Itzykson and J.-B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).

- [6] S. Coleman, *Aspects of Symmetry* (Cambridge University Press, 1985).
- [7] C. Itzykson and J.-M. Drouffe, *Statistical Field Theory* (Cambridge University Press, 1991).
- [8] G. Parisi, *Statistical Field Theory* (Addison Wesley, New York, 1988).
- [9] J. A. Barker and D. Henderson, *Rev. Mod. Phys.* **48**, 587 (1976).
- [10] J. A. Barker, *J. Chem. Phys.* **70**, 2914 (1979).
- [11] M. H. Kalos and P. A. Whitlock, *Monte Carlo Methods*, vol. 1: Basics (John Wiley and Sons, New York, 1986).
- [12] E. L. Pollock and D. M. Ceperley, *Phys. Rev. B* **30**, 2555 (1984).
- [13] D. M. Ceperley, *Rev. Mod. Phys.* **67**, 279 (1995).
- [14] B. Baaquie, *Quantum Finance* (Cambridge University Press, 2004).
- [15] H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets* (World Scientific, 2004).
- [16] M. Takahashi and M. Imada, *J. Phys. Soc. Jpn.* **53**, 3765 (1984).
- [17] X. P. Li and J. Q. Broughton, *J. Chem. Phys.* **86**, 5094 (1987).
- [18] S. Jang, S. Jang, and G. Voth, *J. Chem. Phys.* **115**, 7832 (2001).
- [19] N. Makri and W. H. Miller, *Chem. Phys. Lett.* **151**, 1 (1988).
- [20] N. Makri and W. H. Miller, *J. Chem. Phys.* **90**, 904 (1989).
- [21] N. Makri, *Chem. Phys. Lett.* **193**, 435 (1992).
- [22] M. Alford, T. R. Klassen, and G. P. Lepage, *Phys. Rev. D* **58**, 034503 (1998).
- [23] S. D. Bond, B. B. Laird, and B. J. Leimkuhler, *J. Comp. Phys.* **185**, 472 (2003).
- [24] F. R. Krajewski and M. H. Müser, *Phys. Rev. B* **65**, 174304 (2002).
- [25] A. Erdelyi, *Asyptotic Expansions* (Dover Publications, New York, 1956).
- [26] URL <http://scl.phy.bg.ac.yu/speedup/>.