



C programs for solving the time-dependent Gross–Pitaevskii equation in a fully anisotropic trap[☆]

Dušan Vudragović^a, Ivana Vidanović^a, Antun Balaž^{a,*}, Paulsamy Muruganandam^b, Sadhan K. Adhikari^c

^a Scientific Computing Laboratory, Institute of Physics Belgrade, University of Belgrade, Pregrevica 118, 11080 Belgrade, Serbia

^b School of Physics, Bharathidasan University, Palkalaiperur Campus, Tiruchirappalli – 620024, Tamil Nadu, India

^c Instituto de Física Teórica, UNESP – Universidade Estadual Paulista, 01.140-70 São Paulo, São Paulo, Brazil

ARTICLE INFO

Article history:

Received 2 March 2012

Accepted 21 March 2012

Available online 21 April 2012

Keywords:

Bose–Einstein condensate

Gross–Pitaevskii equation

Split-step Crank–Nicolson scheme

Real- and imaginary-time propagation

C program

OpenMP

Partial differential equation

ABSTRACT

We present C programming language versions of earlier published Fortran programs (Muruganandam and Adhikari (2009) [1]) for calculating both stationary and non-stationary solutions of the time-dependent Gross–Pitaevskii (GP) equation. The GP equation describes the properties of dilute Bose–Einstein condensates at ultra-cold temperatures. C versions of programs use the same algorithms as the Fortran ones, involving real- and imaginary-time propagation based on a split-step Crank–Nicolson method. In a one-space-variable form of the GP equation, we consider the one-dimensional, two-dimensional, circularly-symmetric, and the three-dimensional spherically-symmetric harmonic-oscillator traps. In the two-space-variable form, we consider the GP equation in two-dimensional anisotropic and three-dimensional axially-symmetric traps. The fully-anisotropic three-dimensional GP equation is also considered. In addition to these twelve programs, for six algorithms that involve two and three space variables, we have also developed threaded (OpenMP parallelized) programs, which allow numerical simulations to use all available CPU cores on a computer. All 18 programs are optimized and accompanied by makefiles for several popular C compilers. We present typical results for scalability of threaded codes and demonstrate almost linear speedup obtained with the new programs, allowing a decrease in execution times by an order of magnitude on modern multi-core computers.

New version program summary

Program title: GP-SCL package, consisting of: (i) *imagtime1d*, (ii) *imagtime2d*, (iii) *imagtime2d-th*, (iv) *imagtimecir*, (v) *imagtime3d*, (vi) *imagtime3d-th*, (vii) *imagtimeaxial*, (viii) *imagtimeaxial-th*, (ix) *imagtimesph*, (x) *realttime1d*, (xi) *realttime2d*, (xii) *realttime2d-th*, (xiii) *realttimecir*, (xiv) *realttime3d*, (xv) *realttime3d-th*, (xvi) *realttimeaxial*, (xvii) *realttimeaxial-th*, (xviii) *realttimesph*.

Catalogue identifier: AEDU_v2_0.

Program Summary URL: http://cpc.cs.qub.ac.uk/summaries/AEDU_v2_0.html.

Program obtainable from: CPC Program Library, Queen's University of Belfast, N. Ireland.

Licensing provisions: Standard CPC licence, <http://cpc.cs.qub.ac.uk/licence/licence.html>.

No. of lines in distributed program, including test data, etc.: 180 583.

No. of bytes in distributed program, including test data, etc.: 1 188 688.

Distribution format: tar.gz.

Programming language: C and C/OpenMP.

Computer: Any modern computer with C language compiler installed.

Operating system: Linux, Unix, Mac OS, Windows.

RAM: Memory used with the supplied input files: 2–4 MB (i, iv, ix, x, xiii, xvi, xvii, xviii), 8 MB (xi, xii), 32 MB (vii, viii), 80 MB (ii, iii), 700 MB (xiv, xv), 1.2 GB (v, vi).

Number of processors used: For threaded (OpenMP parallelized) programs, all available CPU cores on the computer.

Classification: 2.9, 4.3, 4.12.

[☆] D.V., I.V., and A.B. acknowledge support by the Ministry of Education and Science of the Republic of Serbia under projects No. ON171017 and NAD-BEC, by DAAD - German Academic and Exchange Service under project NAD-BEC, and by the European Commission under EU FP7 projects PRACE-1IP, PRACE-2IP, HP-SEE, and EGI-InSPIRE. P.M. acknowledges support by DST and CSIR of India. S.K.A. acknowledges support by FAPESP and CNPq of Brazil.

* Corresponding author.

E-mail address: antun@ipb.ac.rs (A. Balaž).

Catalogue identifier of previous version: AEDU_v1_0.

Journal reference of previous version: Comput. Phys. Commun. 180 (2009) 1888.

Does the new version supersede the previous version?: No.

Nature of problem: These programs are designed to solve the time-dependent Gross–Pitaevskii (GP) nonlinear partial differential equation in one-, two- or three-space dimensions with a harmonic, circularly-symmetric, spherically-symmetric, axially-symmetric or fully anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Solution method: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation, in either imaginary or real time, over small time steps. The method yields solutions of stationary and/or non-stationary problems.

Reasons for the new version: Previous Fortran programs [1] are used within the ultra-cold atoms [2–11] and nonlinear optics [12,13] communities, as well as in various other fields [14–16]. This new version represents translation of all programs to the C programming language, which will make it accessible to the wider parts of the corresponding communities. It is well known that numerical simulations of the GP equation in highly experimentally relevant geometries with two or three space variables are computationally very demanding, which presents an obstacle in detailed numerical studies of such systems. For this reason, we have developed threaded (OpenMP parallelized) versions of programs *imagtime2d*, *imagtime3d*, *imagtimeaxial*, *realtime2d*, *realtime3d*, *realtimeaxial*, which are named *imagtime2d-th*, *imagtime3d-th*, *imagtimeaxial-th*, *realtime2d-th*, *realtime3d-th*, *realtimeaxial-th*, respectively. Fig. 1 shows the scalability results obtained for OpenMP versions of programs *realtime2d* and *realtime3d*. As we can see, the speedup is almost linear, and on a computer with the total of 8 CPU cores we observe in Fig. 1(a) a maximal speedup of around 7, or roughly 90% of the ideal speedup, while on a computer with 12 CPU cores we find in Fig. 1(b) that the maximal speedup is around 9.6, or 80% of the ideal speedup. Such a speedup represents significant improvement in the performance.

Summary of revisions: All Fortran programs from the previous version [1] are translated to C and named in the same way. The structure of all programs is identical. We have introduced the use of comprehensive input files, where all parameters are explained in detail and can be set by a user. We have also included makefiles with tested and verified settings for GNU's gcc compiler, Intel's icc compiler, IBM's xlc compiler, PGI's pgcc compiler, and Oracle's suncc (former Sun's) compiler. In addition to this, 6 new threaded (OpenMP parallelized) programs are supplied (*imagtime2d-th*, *imagtime3d-th*, *imagtimeaxial-th*, *realtime2d-th*, *realtime3d-th*, *realtimeaxial-th*) for algorithms involving two or three space variables. They are written by OpenMP-parallelizing the most computationally demanding loops in functions performing time evolution (*calcnu*, *calclux*, *calcluy*, *calcluz*), normalization (*calcnorm*), and calculation of physical quantities (*calcmuen*, *calcrms*). Since some of the dynamically allocated array variables are used within such loops, they had to be made private for each thread. This was done by allocating matrices instead of arrays, with the first index in all such matrices corresponding to a thread number.

Additional comments: This package consists of 18 programs, see Program title above, out of which 12 programs (i, ii, iv, v, vii, ix, x, xi, xiii, xiv, xvi, xviii) are serial, while 6 programs (iii, vi, viii, xii, xv, xvii) are threaded (OpenMP parallelized). For the particular purpose of each program, please see descriptions below.

Running time: All running times given in descriptions below refer to programs compiled with gcc on quad-core Intel Xeon X5460 at 3.16 GHz (CPU1), and programs compiled with icc on quad-core Intel Nehalem E5540 at 2.53 GHz (CPU2). With the supplied input files, running times on CPU1 are: 5 min (i, iv, ix, xii, xiii, xvii, xviii), 10 min (viii, xvi), 15 min (iii, x, xi), 30 min (ii, vi, vii), 2 h (v), 4 h (xv), 15 h (xiv). On CPU2, running times are: 5 min (i, iii, iv, viii, ix, xii, xiii, xvi, xvii, xviii), 10 min (vi, x, xi), 20 min (ii, vii), 1 h (v), 2 h (xv), 12 h (xiv).

© 2012 Elsevier B.V. All rights reserved.

New version program summary (i)

Program title: *imagtime1d*.

Title of electronic files: *imagtime1d.c*, *imagtime1d.h*.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 4 MB.

Programming language used: C.

Typical running time: 2 min (CPU1), 1 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in one space dimension with a harmonic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (ii)

Program title: *imagtime2d*.

Title of electronic files: *imagtime2d.c*, *imagtime2d.h*.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 80 MB.

Programming language used: C.

Typical running time: 30 min (CPU1), 20 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in two space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (iii)

Program title: imagtime2d-th.

Title of electronic files: imagtime2d-th.c, imagtime2d-th.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 80 MB.

Programming language used: C/OpenMP.

Typical running time: 15 min (CPU1), 5 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in two space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (iv)

Program title: imagtimecir.

Title of electronic files: imagtimecir.c, imagtimecir.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 2 MB.

Programming language used: C.

Typical running time: 2 min (CPU1), 1 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in two space dimensions with a circularly-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (v)

Program title: imagtime3d.

Title of electronic files: imagtime3d.c, imagtime3d.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 1.2 GB.

Programming language used: C.

Typical running time: 1.5 h (CPU1), 1 h (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (vi)

Program title: imagtime3d-th.

Title of electronic files: imagtime3d-th.c, imagtime3d-th.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 1.2 GB.

Programming language used: C/OpenMP.

Typical running time: 25 min (CPU1), 10 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation

in three space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (vii)

Program title: imagtimeaxial.

Title of electronic files: imagtimeaxial.c, imagtimeaxial.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 32 MB.

Programming language used: C.

Typical running time: 30 min (CPU1), 20 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with an axially-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (viii)

Program title: imagtimeaxial-th.

Title of electronic files: imagtimeaxial-th.c, imagtimeaxial-th.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 32 MB.

Programming language used: C/OpenMP.

Typical running time: 10 min (CPU1), 5 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with an axially-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (ix)

Program title: imagtimesph.

Title of electronic files: imagtimesph.c, imagtimesph.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 2.5 MB.

Programming language used: C.

Typical running time: 2 min (CPU1), 1 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with a spherically-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in imaginary time over small time steps. The method yields solutions of stationary problems.

New version program summary (x)

Program title: realtime1d.

Title of electronic files: realtime1d.c, realtime1d.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 4 MB.

Programming language used: C.

Typical running time: 15 min (CPU1), 10 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in one space dimension with a harmonic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

New version program summary (xi)

Program title: realtime2d.

Title of electronic files: realtime2d.c, realtime2d.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 8 MB.

Programming language used: C.

Typical running time: 15 min (CPU1), 10 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in two space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

New version program summary (xii)

Program title: realtime2d-th.

Title of electronic files: realtime2d-th.c, realtime2d-th.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 8 MB.

Programming language used: C/OpenMP.

Typical running time: 5 min (CPU1), 2 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in two space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

New version program summary (xiii)

Program title: realtimecir.

Title of electronic files: realtimecir.c, realtimecir.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 3 MB.

Programming language used: C.

Typical running time: 5 min (CPU1), 5 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in two space dimensions with a circularly-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

New version program summary (xiv)

Program title: realtime3d.

Title of electronic files: realtime3d.c, realtime3d.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 700 MB.

Programming language used: C.

Typical running time: 15 h (CPU1), 12 h (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

New version program summary (xv)

Program title: realtime3d-th.

Title of electronic files: realtime3d-th.c, realtime3d-th.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 700 MB.

Programming language used: C/OpenMP.

Typical running time: 4 h (CPU1), 1.8 h (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with an anisotropic trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

New version program summary (xvi)

Program title: realtimeaxial.

Title of electronic files: realtimeaxial.c, realtimeaxial.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 4 MB.

Programming language used: C.

Typical running time: 10 min (CPU1), 5 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with an axially-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.




Fig. 1. (Colour online) Speedup in the execution time of realtime2d-th and realtime3d-th threaded (OpenMP parallelized) programs as a function of the number of CPU cores used. The results are obtained: (a) on an 8-core machine with $2 \times$ quad-core Intel Nehalem E5540 CPU at 2.53 GHz, using the icc compiler, (b) on a 12-core machine with $2 \times$ six-core Intel Nehalem X5650 CPU at 2.66 GHz, using the pgcc compiler. The spatial grid sizes used are 2000×2000 (realtime2d-th) and $1000 \times 1000 \times 300$ (realtime3d-th).

New version program summary (xvii)

Program title: realtimeaxial-th.

Title of electronic files: realtimeaxial-th.c, realtimeaxial-th.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 4 MB.

Programming language used: C/OpenMP.

Typical running time: 5 min (CPU1), 1 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with an axially-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

New version program summary (xviii)

Program title: reallimesph.

Title of electronic files: reallimesph.c, reallimesph.h.

Computer: Any modern computer with C language compiler installed.

Maximum RAM memory: 2.5 MB.

Programming language used: C.

Typical running time: 5 min (CPU1), 5 min (CPU2).

Nature of physical problem: This program is designed to solve the time-dependent GP nonlinear partial differential equation in three space dimensions with a spherically-symmetric trap. The GP equation describes the properties of a dilute trapped Bose–Einstein condensate.

Method of solution: The time-dependent GP equation is solved by the split-step Crank–Nicolson method by discretizing in space and time. The discretized equation is then solved by propagation in real time over small time steps. The method yields solutions of stationary and non-stationary problems.

References

- [1] P. Muruganandam, S.K. Adhikari, Fortran programs for the time-dependent Gross–Pitaevskii equation in a fully anisotropic trap, *Comput. Phys. Commun.* 180 (2009) 1888.
- [2] G. Mazzarella, L. Salasnich, Collapse of triaxial bright solitons in atomic Bose–Einstein condensates, *Phys. Lett. A* 373 (2009) 4434.
- [3] Y. Cheng, S.K. Adhikari, Symmetry breaking in a localized interacting binary Bose–Einstein condensate in a bichromatic optical lattice, *Phys. Rev. A* 81 (2010) 023620; S.K. Adhikari, H. Lu, H. Pu, Self-trapping of a Fermi superfluid in a double-well potential in the Bose–Einstein-condensate-unity crossover, *Phys. Rev. A* 80 (2009) 063607.
- [4] S. Gautam, D. Angom, Rayleigh–Taylor instability in binary condensates, *Phys. Rev. A* 81 (2010) 053616; S. Gautam, D. Angom, Ground state geometry of binary condensates in axisymmetric traps, *J. Phys. B* 43 (2010) 095302; S. Gautam, P. Muruganandam, D. Angom, Position swapping and pinching in Bose–Fermi mixtures with two-color optical Feshbach resonances, *Phys. Rev. A* 83 (2011) 023605.
- [5] S.K. Adhikari, B.A. Malomed, L. Salasnich, F. Toigo, Spontaneous symmetry breaking of Bose–Fermi mixtures in double-well potentials, *Phys. Rev. A* 81 (2010) 053630.
- [6] G.K. Chaudhary, R. Ramakumar, Collapse dynamics of a $(176)\text{Yb}-(174)\text{Yb}$ Bose–Einstein condensate, *Phys. Rev. A* 81 (2010) 063603.
- [7] S. Sabari, R.V.J. Raja, K. Porsezian, P. Muruganandam, Stability of trapless Bose–Einstein condensates with two- and three-body interactions, *J. Phys. B* 43 (2010) 125302.
- [8] L.E. Young-S, L. Salasnich, S.K. Adhikari, Dimensional reduction of a binary Bose–Einstein condensate in mixed dimensions, *Phys. Rev. A* 82 (2010) 053601.
- [9] I. Vidanović, A. Balaž, H. Al-Jibbouri, A. Pelster, Nonlinear Bose–Einstein-condensate dynamics induced by a harmonic modulation of the s -wave scattering length, *Phys. Rev. A* 84 (2011) 013618.
- [10] R.R. Sakhel, A.R. Sakhel, H.B. Ghassib, Self-interfering matter-wave patterns generated by a moving laser obstacle in a two-dimensional Bose–Einstein condensate inside a power trap cut off by box potential boundaries, *Phys. Rev. A* 84 (2011) 033634.
- [11] A. Balaž, A.I. Nicolin, Faraday waves in binary nonmiscible Bose–Einstein condensates, *Phys. Rev. A* 85 (2012) 023613; A.I. Nicolin, Variational treatment of Faraday waves in inhomogeneous Bose–Einstein condensates, *Physica A* 391 (2012) 1062; A.I. Nicolin, Resonant wave formation in Bose–Einstein condensates, *Phys. Rev. E* 84 (2011) 056202; A.I. Nicolin, Faraday waves in Bose–Einstein condensates subject to anisotropic transverse confinement, *Rom. Rep. Phys.* 63 (2011) 1329; A.I. Nicolin, M.C. Raportaru, Faraday waves in high-density cigar-shaped Bose–Einstein condensates, *Physica A* 389 (2010) 4663.
- [12] S. Yang, M. Al-Amri, J. Evers, M.S. Zubairy, Controllable optical switch using a Bose–Einstein condensate in an optical cavity, *Phys. Rev. A* 83 (2011) 053821.
- [13] W. Hua, X.-S. Liu, Dynamics of cubic and quintic nonlinear Schrödinger equations, *Acta Phys. Sinica* 60 (2011) 110210.
- [14] Z. Sun, W. Yang, An exact short-time solver for the time-dependent Schrödinger equation, *J. Chem. Phys.* 134 (2011) 041101.
- [15] A. Balaž, I. Vidanović, A. Bogojević, A. Belić, A. Pelster, Fast converging path integrals for time-dependent potentials: I. Recursive calculation of short-time expansion of the propagator, *J. Stat. Mech. Theory Exp.* (2011) P03004.
- [16] W.B. Cardoso, A.T. Avelar, D. Bazeia, One-dimensional reduction of the three-dimensional Gross–Pitaevskii equation with two- and three-body interactions, *Phys. Rev. E* 83 (2011) 036604.